

TOM J. POUSTI, MD, F.A.C.S.
PLASTIC AND RECONSTRUCTION SURGERY

INFORMED-CONSENT-CAPSULECTOMY WITH BREAST IMPLANT EXCHANGE

INSTRUCTIONS

This is an informed-consent document that has been prepared to help inform you about open capsulectomy with breast implant exchange, its risks, and alternative treatments.

It is important that you read this information carefully and completely. Please initial each page, indicating that you have read the page and sign the consent for surgery as proposed by your plastic surgeon.

GENERAL INFORMATION

Open capsulectomy with breast implant exchange is a surgical operation performed to treat scarring with occurs around breast implants and to place new breast implants. Individuals with old, damaged or broken implants may consider open capsulectomy surgery with breast implant exchange as a procedure to maintain the long-term results from their original surgery. You may be advised by your surgeon to consider replacing your breast implants with new ones, irrespective of how long you have had them. In some situations, you may have been advised to consider new breast implants with a textured outer surface. Breast implants do not have an indefinite life span and will eventually require replacement surgery.

Scar tissue, which forms internally around a breast implant, can tighten and make the breast round, firm, and possibly painful. Excessive firmness of the breasts can occur soon after the original surgery or years later. The incidence of symptomatic capsular contracture can be expected to increase over time. Capsular contracture may occur on one side, both sides or not at all. Calcification can occur within the scar tissue that surrounds breast implants. Treatment for capsular contracture may require surgery, implant replacement, or implant removal.

Depending on the extent of the scarring problem, it may be necessary to place the implant in a deeper location, underneath the pectoralis muscle on the chest. Incisions for the open capsulectomy procedure may be placed in different locations than those used for the original surgery. If the breasts are not the same size or shape before surgery, it is unlikely that they will be completely symmetrical afterward.

Patients undergoing open capsulectomy surgery with breast implant exchange must consider the possibility of future revisionary surgery. Breast implants cannot be expected to last forever.

ALTERNATIVE TREATMENT

Open capsulectomy with breast implant exchange is an elective surgical operation. Alternative treatment would consist of removal of your breast implants. Risks and potential complications are associated with alternative treatments that involve surgery.

RISKS OF OPEN CAPSULECTOMEY WITH BREAST IMPLANT EXCHANGE

Every surgical procedure involves a certain amount of risk and it is important that you understand the risks involved with open capsulectomy with breast implant exchange. Additional information concerning breast implants may be obtained from the FDA, package-insert sheets supplied by the implant manufacturer, or other information pamphlets required by the individual state laws.

An individual's choice to undergo a surgical procedure is based on the comparison of the risk to potential benefit. Although the majority of women do not experience the following complications, you should discuss each of them with your plastic surgeon to make sure you understand the risks, potential complications, and consequences of open capsulectomy with breast implant exchange.

BLEEDING

It is possible, though unusual, to experience a bleeding episode during or after surgery. Should post-operative bleeding occur, it may require emergency treatment to drain accumulated blood (hematoma). Do not take any aspirin or anti-inflammatory medications for ten days before surgery, as this may increase the risk of bleeding.

INFECTION

Infection is unusual after this type of surgery. It may appear in the immediate post-operative period or at any time following the insertion of a breast implant. Sub-acute or chronic infections may be difficult to diagnose. Should an infection occur, treatment including antibiotics, possible removal of the implant, or additional surgery may be necessary. Infections with the presence of a breast implant are harder to treat than infections in normal body tissues. If an infection does not respond to antibiotics, the breast implant may have to be removed. After the infection is treated, a new breast implant can usually be reinserted. It is extremely rare that an infection would occur around an implant from a bacterial infection elsewhere in the body, however, prophylactic antibiotics may be considered for subsequent dental or other surgical procedures.

RECURRENT CAPSULAR CONTRACTURE

Excessive firmness of the breasts can re-occur soon after capsulectomy surgery or years later. Scar tissue may form again around the breast implant; it can tighten and make the breast round, firm, and possibly painful. The recurrence of symptomatic capsular contracture following capsulectomy is not predictable. The incidence of symptomatic capsular contracture can be expected to increase over time. Capsular contracture may re-occur on one side, both sides or not at all. Treatment for recurrent capsular contracture may require additional surgery, or implant removal.

INABILITY TO REMOVE CAPSULAR SCAR TISSUE

It may not be possible to completely remove all scar tissue that has formed around the breast implant(s). It may not be possible to completely remove textured implant shell material that has become adherent to the scar layer.

DAMAGE TO ADJACENT STRUCTURES

Scar tissue may be adherent to breast tissue, muscle or the chest wall. These structures may be damaged during capsulectomy surgery to remove the scar capsule.

ESCAPED SILICONE GEL MATERIAL

In situations that involve damaged, leaking, or ruptured silicone gel breast implants, gel material can escape into the space inside the scar capsule layer around the breast implant. In other situations, gel material may have escaped outside of the scar capsule layer into the breast and other tissue. Additional surgical procedures may be needed to remove escaped gel material. It may not be possible to completely remove escaped gel material discovered during surgery.

CHANGE IN NIPPLE AND SKIN SENSATION

Some change in nipple sensation is not unusual right after surgery. After several months, most patients have normal sensation. Partial or permanent loss of nipple and skin sensation may occur occasionally.

SKIN SCARRING

Excessive scarring is uncommon. In rare cases, abnormal scars may result. Scars may be unattractive and of different color than surrounding skin. Additional surgery may be needed to treat abnormal scarring after surgery.

IMPLANTS

Breast implants, similar to other medical devices, can fail. Implants can break or leak. When a saline-filled implant deflates, its salt water filling will be absorbed by the body. Rupture can occur as a result of an injury, from no apparent cause, or during a mammography. It is possible to damage an implant at the time of surgery. Damaged or broken implants cannot be repaired. Ruptured or deflated implants require replacement or removal. Breast implants cannot be expected to last forever.

DEGRADATION OF BREAST IMPLANTS

It is possible that small pieces of the implant material may separate from the outer surface of breast implants. This is of unknown significance.

SURFACE CONTAMINATION OF IMPLANTS

Skin oil, lint from surgical drapes, or talc may become deposited on the surface of the implant at the time of insertion. The consequences of this are unknown.

IMPLANT EXTRUSION

Lack of adequate tissue coverage or infection may result in exposure and extrusion of the implant. Skin breakdown has been reported with the use of steroid drugs or after radiation therapy to breast tissue. If tissue breakdown occurs and the implant becomes exposed, implant removal may be necessary. Smoking may interfere with the healing process.

CALCIFICATION

Calcium deposits can form in the scar tissue surrounding the implant and may cause pain, firmness, and be visible on mammography. These deposits must be identified as different from calcium deposits that are a sign of breast cancer. Should this occur, additional surgery may be necessary to remove and examine calcifications.

SKIN WRINKLING AND RIPPLING

Visible and palpable wrinkling of implants can occur. Some wrinkling is normal and expected. This may be more pronounced in patients who have saline filled implants or thin breast tissue. It may be possible to feel the implant valve. Some patients may find palpable valve and wrinkles cosmetically undesirable. Palpable valve, wrinkling and/or folds may be confused with palpable tumors and questionable cases must be investigated. An implant may become visible at the surface of the breast as a result of the device pushing through the layers of skin.

MAMMOGRAPHY

Breast implants may make mammography more difficult and may obscure the detection of breast cancer. Implant rupture can occur from breast compression during mammography. Inform your mammography technologist of the presence of breast implants so that appropriate mammogram studies may be obtained. Patients with capsular contracture may find mammogram techniques painful and the difficulty of breast imaging will increase with the extent of contracture. Ultrasound, specialized mammography and MRI studies may be of benefit to evaluate breast lumps and the condition of the implant(s). Because more x-ray views are necessary with specialized mammography techniques, women with breast implants will receive more radiation than women without implants who receive a normal exam. However, the benefit of the mammogram in finding cancer outweighs the risk of additional x-rays.

PREGNANCY AND BREAST FEEDING

Although many women with breast implants have successfully breast fed their babies, it is not known if there are increased risks in nursing for a woman with breast implants or if the children of women with breast implants are more likely to have health problems. There is insufficient evidence regarding the absolute safety of breast implants in relation of fertility, pregnancy or breast feeding. Some women with breast implants have reported health problems in the breast fed children. Only very limited research has been conducted in this area and at this time there is no scientific evidence that this is a problem.

IMPLANT DISPLACEMENT

Displacement or migration of a breast implant may occur from its initial placement and can be accompanied by discomfort and/or distortion in breast shape. Difficult techniques of implant placement may increase the risk of displacement or migration.

SURGICAL ANESTHESIA

Both local and general anesthesia involves risk. There is the possibility of complications, injury, and even death from all forms of surgical anesthesia or sedation.

CHEST WALL DEFORMITY

Chest wall deformity has been reported secondary to the use of tissue expanders and breast implants. The consequences of chest wall deformity are of unknown significance.

UNUSUAL ACTIVITIES AND OCCUPATIONS

Activities and occupations which have the potential for trauma to the breast could potentially break or damage breast implants, or cause bleeding.

ALLERGIC REACTIONS

In rare cases, local allergies to tape, suture material, or topical preparations have been reported. Systemic reactions, which are more serious, may result from drugs used during surgery and prescription medicines. Allergic reactions may require additional treatment.

SEROMA

Fluid may accumulate around the implant following surgery, trauma or vigorous exercise. Additional treatment may be necessary to drain fluid accumulation around breast implants.

BREAST DISEASE

Current medical information does not demonstrate an increased risk of breast disease or breast cancer in women who have breast implant surgery for either cosmetic or reconstructive purposes. Breast disease can occur independently of breast implants. It is recommended that all women perform periodic self examination of their breasts, have a mammography according to American Cancer Society guidelines, and seek professional care should the notice a breast lump.

LONG TERM RESULTS

Subsequent alterations in breast shape may occur as the result of aging, weight loss or gain, pregnancy, or other circumstances not related to open capsulectomy with breast implant exchange. Breast sagging may normally occur.

THROMBOSED VEINS

Thrombosed veins, which resemble cords, occasionally develop in the area of the breast and resolve without medical or surgical treatment.

IMMUNE SYSTEM DISEASES AND UNKNOWN RISKS

Some women with breast implants have reported symptoms similar to those of known diseases of the immune system, such as systemic lupus erthematosus, rheumatoid arthritis, scleroderma, and other arthritis like conditions. A connection between implanted silicone and connective tissue disorders has been reported in the medical literature. To date, there is no scientific evidence that women with either silicone gel-filled or saline-filled breast implants have an increased risk of these diseases, but the possibility cannot be excluded. If a casual relationship is established, the theoretical risk of immune and unknown disorders may be low. The effects of breast implants in individuals with pre-existing connective tissue disorders is unknown.

Unlike silicone gel-filled implants, the saline-filled implants contain salt water. Any risk related to silicone gel would not be associated with saline-filled implants. However, gel-filled and saline-filled devices have a silicone rubber envelope. An increased risk of auto-immune disease is possible even from saline implants. Reliable medical laboratory tests have determined antibodies to silicone do not exist. It has not been proved that there is a relationship between silicone antibodies and disease in women with breast implants. Currently, there is insufficient evidence to state that there is a health benefit from removing either breast implant(s) and scar tissue capsule(s) or that removal will alter auto-immune disease or prevent its potential occurrence.

In very few women who have breast implants, a variety of other symptoms and conditions have been reported, suggestive of an auto-immune multiple-sclerosis like syndrome. Additional complaints involve the muscular skeletal, skin, nervous, and immune systems. The relationship of breast implants to these conditions has been hypothesized, although not scientifically proven. Because such disease states are rare, they are difficult to research.

Current studies have only looked for the symptoms of known auto-immune diseases, rather than the variety of symptoms that women report experiencing. Some of the reported symptoms include:

- ◆ Swelling and/or joint pain or arthritis like pain
- ◆ General aching
- ◆ Unusual hair loss
- ◆ Unexplained or unusual; loss of energy
- ◆ Greater chance of getting colds, viruses, flu
- ◆ Swollen glands or lymph nodes
- ◆ Rash, fever
- ◆ Memory problems, headaches
- ◆ Muscle weakness or burning
- ◆ Nausea, vomiting
- ◆ Irritable bowel syndrome

Questions have been raised about the potential for the saline solution used to fill breast implants to become contaminated with bacteria or fungus. These organisms may present a risk to the patient in the event of implant leakage or deflation.

There is the possibility of unknown risks associated with silicone breast implants and tissue expanders

TOXIC SHOCK SYNDROME

This is an extremely rare complication following breast augmentation, reconstruction, or tissue expansion with silicone implants.

UNSATISFACTORY RESULT

You may be disappointed with the results of surgery. Asymmetry in implant placement, breast shape and size may occur after surgery. Unsatisfactory surgical scar location or displacement may occur. Pain may occur following surgery. It may be necessary to perform additional surgery to improve your results.

REMOVAL/REPLACEMENT OF BREAST IMPLANTS

Future removal or replacement of breast implants and the surrounding scar tissue envelope involves a surgical procedure with risks and potential complications.

HEALTH INSURANCE

Most health insurance companies exclude coverage for cosmetic surgical operations and any complications that might occur from surgery. Some insurance carriers may possibly exclude breast diseases in patients who have breast implants. However, in some situations involving breast implants placed for the purpose of breast reconstruction, your insurance plan may consider open capsulectomy with implant exchange a covered benefit. Please carefully review your health insurance subscriber information pamphlet and call your insurance company.

ADDITIONAL SURGERY NECESSARY

Should complications occur, additional surgery or other treatments may be necessary. Even though risks and complications occur infrequently, the risks cited are particularly associated with open capsulectomy with breast implant exchange; other complications and risks can occur but are even more uncommon. The practice of medicine and surgery is not an exact science. Although good results are expected, there is no guarantee or warranty expressed or implied on the results that may be obtained.

FINANCIAL RESPONSIBILITIES

The cost of surgery involves several charges for the services provided. The total includes fees charged by your doctor, the cost of implants and surgical supplies, anesthesia, laboratory tests, and possible out patient hospital charges, depending on where the surgery is performed. Depending on whether the cost of surgery is covered by an insurance plan, you will be responsible for necessary co-payments, deductibles, and charges not covered. Additional costs may occur should complications develop from the surgery. Secondary surgery or hospital day surgery charges involved with revisionary surgery would also be your responsibility.

DISCLAIMER

Informed-consent documents are used to communicate information about the proposed surgical treatment of a disease or condition along with disclosure of risks and alternative forms of treatment(s). The informed consent process attempts to define principles or risk disclosure that should generally meet the needs of most patients in most circumstances.

However, informed consent documents should not be considered all inclusive in defining other methods of care and risks encountered. Your plastic surgeon may provide you with additional or different information, which is based on all the facts in your particular case and the state of medical knowledge.

Informed consent documents are not intended to define or serve as the standard of medical care. Standards of medical care are determined on the basis of all the facts involved in an individual case and are subject to change as scientific knowledge and technology advance and as practice patterns evolve.

It is important that you read the above information carefully and have all of your questions answered before signing the consent on the next page.

CONSENT FOR SURGERY/PROCEDURE or TREATMENT

I hereby authorize Dr. Tom J. Pousti and such assistant as may be selected to perform the following procedure or treatment:

CAPSULECTOMY SURGERY

I have received the following information sheet:

INFORMED CONSENT FOR CAPSULECTOMY WITH IMPLANT EXCHANGE

I recognize that during the course of the operation and medical treatment or anesthesia, unforeseen conditions may necessitate different procedures than those above. I therefore authorize the above physician and assistants or designees to perform such other procedures that are in the exercise of his or her professional judgment necessary and desirable. The authority granted under this paragraph shall include all conditions that require treatment and are not known to my physician at the time the procedure has begun.

I consent to the administration of such anesthetics considered necessary or advisable. I understand that all forms of anesthesia involve risk and possibility of complications, injury, and sometimes death.

I acknowledge that no guarantee has been given by anyone as to the results that may be obtained.

I consent to the photographing or televising of the operation or procedure to be performed, including appropriate portions of my body, for medical, scientific or educational purposes, provided my identity is not revealed by the pictures.

For purposes of advancing medical education, I consent to the admittance of observers to the operating room.

I consent to the disposal of any tissue, medical devices or body parts, which may be removed.

I authorize the release of my social security number to appropriate agencies for legal reporting and medical device registration if applicable.

IT HAS BEEN EXPLAINED TO ME IN A WAY THAT I UNDERSTAND:

- a) THE ABOVE TREATMENT OR PROCEDURE TO BE UNDERTAKEN
- b) THERE MAY BE ALTERNATIVE PROCEDURE OR METHODS OF TREATMENT
- c) THERE ARE RISKS TO THE PROCEDURE OR TREATMENT PROPOSED

Sign A or B Below

- A. I consent to the treatment or procedure and the above listed items. I have been asked if I want more detailed explanation, but I am satisfied with the explanation and do not want more information.

Patient or person authorized to sign for patient

Date

- B. I consent to the treatment or procedure and above listed items. I requested and received, in substantial detail, further explanation of the procedure or treatment, other alternative procedures or methods or treatment and information about the material risks or the procedure of treatment.

Patient or person authorized to sign for patient

Date